

Emergence and control of an outbreak of infections due to Panton-Valentine leukocidin positive, ST22 methicillin-resistant *Staphylococcus aureus* in a neonatal intensive care unit

A. N. Pinto¹, R. Seth², F. Zhou¹, J. Tallon⁴, K. Dempsey⁴, M. Tracy^{2,5}, G. L. Gilbert^{1,3,4} and M. V. N. O'Sullivan^{1,3}

1) Centre for Infectious Diseases and Microbiology, Westmead Hospital, Sydney, 2) Centre for Newborn Care, Westmead Hospital, Sydney,

3) Sydney Emerging Infections and Biosecurity Institute, University of Sydney, 4) Infection Prevention and Control Unit, Westmead Hospital, Sydney and

5) Department of Paediatrics and Child Health, University of Sydney, Sydney, Australia

Abstract

Methicillin resistant *Staphylococcus aureus* (MRSA) infection can cause significant morbidity and mortality in neonates. We investigated a nosocomial MRSA outbreak in a neonatal intensive care unit (NICU), using a novel typing method. Following two fatal cases, in May 2011, a prospective outbreak investigation was conducted, involving neonates, mothers and healthcare workers in a large tertiary NICU in Sydney. MRSA isolates were characterized by antimicrobial susceptibility testing, a multiplex PCR-based reverse line blot (mPCR/RLB) binary typing system and other molecular typing methods. Over 7 months, 14 neonates were colonized with MRSA and six infected: three with superficial lesions and three with life-threatening disease, including the two index cases, who died despite empirical treatment with vancomycin. Isolates from 15 neonates were indistinguishable by RLB typing and identified as a PVL-producing ST22 SCCmec IV MRSA strain, which was resistant to gentamicin and trimethoprim-sulphamethoxazole. The outbreak strain was also isolated from one healthcare worker, one environmental swab and one father, but the source remained obscure. During the same period several different non-multiresistant and multiresistant MRSA strains were isolated from five neonates, five mothers (including two whose infants were colonized with the outbreak strain), one father, three healthcare workers and two environmental swabs. Rapid turnaround time of typing results allowed us to recognize and define the outbreak and implement targeted infection control interventions. PVL-producing ST22 SCCmec IV MRSA appears to be a virulent and highly transmissible pathogen in the NICU, which was difficult to control.

Keywords: Control, MRSA, NICU, outbreak, typing

Original Submission: 19 April 2012; **Revised Submission:** 3 July 2012; **Accepted:** 3 July 2012

Editor: G. Lina

Article published online: 26 July 2012

Clin Microbiol Infect 2013; **19**: 620–627

10.1111/j.1469-0691.2012.03987.x

Corresponding author: Dr A. N. Pinto, Dr M. O'Sullivan, Centre for Infectious Diseases and Microbiology Laboratory Services, ICPMR Building, Level 3, Westmead Hospital, Darcy Road, Westmead NSW 2145, Australia

E-mails: pinto.angie@gmail.com; matthew.osullivan@swahs.health.nsw.gov.au

This is the first description of an outbreak due to a rare clone of MRSA using a novel typing system. This highly discriminatory method is inexpensive, high throughput, has a turnaround time of 12 h, and enabled targeted infection control interventions.

Introduction

Methicillin-resistant *Staphylococcus aureus* (MRSA) has the potential to cause severe disease and sustained nosocomial

outbreaks [1–3] among premature infants in neonatal intensive care units (NICUs). Its early recognition is essential to prevent transmission and ensure optimal treatment [4].

The mother is often assumed to be the source of MRSA in a colonized neonate, but nosocomial transmission from fomites or via healthcare workers (HCWs) also occurs. Transmission routes can be identified with the aid of molecular typing [3].

In May 2011, two extremely premature (26 weeks gestation) infants developed fulminant MRSA sepsis and died within a few days of each other. This paper describes the outbreak investigation, using a novel rapid MRSA strain-typing method.

Methods

Design

This was a prospective study in which infection control interventions were based upon molecular typing data available in real time.

Setting

Westmead Hospital is a 980-bed teaching hospital in Sydney, Australia, serving a population of 1.5 million; there are c. 5500 deliveries annually. The 39-bed NICU comprises 19 high acuity ventilator cots and 20 lower acuity special care cots; in 2011 there were 1635 admissions, including over 100 infants <1500 grams birth weight; 1066 patients were admitted during the outbreak period (18 May 2011 to 23 January 2012). The standard nurse/bed ratio is 1:1–1:4, depending upon staffing levels and patient acuity. There are no single cot isolation rooms.

Patients and case definition

Outbreak cases were defined by isolation of the MRSA outbreak strain from any culture during the outbreak period. In these infants, infection was defined by clinical and laboratory criteria and a requirement for antimicrobial therapy, and colonization by absence of relevant symptoms.

Interventions

MRSA surveillance. Previously, there had been no routine MRSA screening in the NICU because MRSA infections were rare. After two index cases were identified (Fig. 1), all neonates and NICU staff were screened and, subsequently, neonates were swabbed on admission and weekly thereafter. Because of ongoing MRSA acquisitions, NICU staff screening was repeated and extended to delivery room staff in September 2011. Swabs were collected from the anterior nares, umbilicus (if moist) and perineum of neonates and anterior nares, only, of staff. Mothers of colonized or infected neo-

nates were screened using nose, throat, perineal and low vaginal swabs. Frequently touched environmental surfaces, including door handles, trolleys, computer keyboards, ultrasound probes and arterial blood gas and X ray machines, were swabbed in May and September 2011, using sterile, dry swabs, which were inoculated directly into broth (see below).

Decolonization regimen. MRSA-colonized HCWs were treated with mupirocin 2% nasal ointment three times a day and daily triclosan 1% body wash for 5 days.

Culture and strain typing

Surveillance swabs from individual infants and mothers were inoculated, together, into a single tube of methicillin, aztreonam, mannitol salt (MAMS) broth. Environmental swabs were inoculated directly into MAMS broth. Broths were incubated for 18 to 24 h at 35°C and subcultured onto chromogenic Brilliance™ MRSA selective media (Oxoid, Basingstoke, Hants, UK). Molecular methods were also employed for rapid detection, using the MRSA4 Easy-Plex assay kit (AusDiagnostics, Sydney, Australia), which targeted *nuc*, *mecA* and *SCCmec-orfX*; all positive samples were confirmed by culture. Pure cultures were stored in nutrient broth with 20% glycerol at –70°C. Isolates were identified by routine laboratory methods [5], including antibiotic susceptibility testing performed by BD Phoenix Automated Microbiology System (Becton Dickinson, Sparks, MD, USA).

Molecular typing studies

Isolates were typed using a novel binary typing system for MRSA (O'Sullivan MV, Sintchenko V, Gilbert GL; personal communication). The 19 targets for this assay were chosen from 51 utilized in previous assays for toxin gene [6], phage-derived open reading frame [7] and *SCCmec* typing [8]. Results of these assays, using a diverse local and international collection of 165 MRSA isolates, were analysed by a specially designed computerized algorithm (AuSeTTS, available at




FIG. 1. Chest radiographs of Cases 1 and 2. (a) Case 1: rapidly progressive necrotizing pneumonia. (b) Case 2: overwhelming sepsis.

<http://www.cidmph.org.au/pages/AuSeTTS>) (O'Sullivan MV, Sintchenko V, Gilbert GL; submitted for publication) to identify the combination of targets with the highest discriminatory power, while maintaining concordance with multilocus sequence typing (MLST). Targets selected were four toxin genes (*sea*, *sec*, *sed* and *lukS-PV*), nine derived from integrated prophages (Tn554tnpB, ϕ Mu50B SAV0881, ϕ PV83 ORF 2, ϕ Mu50B SAV0858, ϕ 11 nt 4427-5251, ϕ SLT ORF 257, ϕ N315 SA1801, ϕ Mu50A SAV1974 and ϕ SLT ORF 182) and six SCCmec elements (*ccrAB*, *ccrC*, *mecRI*, E007, CQ002 and *cadB*) plus nuc (*S. aureus* control) and *mecA* (MRSA control). All targets were amplified in a single multiplex PCR reaction; products were hybridized to probes on a reusable nylon membrane and detected by chemiluminescence, visualized on X-ray film [9]. Results were expressed as a 19-digit binary number, converted to a decimal code for ease of interpretation. The method's discriminatory power (Simpson's index of diversity, $D = 0.994$; 95% CI, 0.988–1.00) was similar to that of PFGE ($D = 0.987$; 95% CI, 0.977–0.998) and higher than that of *spa* typing ($D = 0.926$; 95% CI, 0.879–0.972) and its concordance with MLST was high (adjusted Wallace coefficient 0.993; 95% CI, 0.986–1.000 [10]). The assay was performed weekly and, during the outbreak investigation, selected isolates from the NICU were further characterized.

Pulse field gel electrophoresis (PFGE) was performed according to the harmony protocol [11]; patterns were examined using BIONUMERICS v3.00 software (Applied Maths, Sint-Martens-Latem, Belgium). PFGE types were defined by

indistinguishable patterns (100% similarity). *spa* typing [6] and MLST [12] were performed according to previously published methods. SCCmec subtyping was performed using mPCR/RLB assay [8]; subtypes were assigned as proposed by Chongtrakool *et al.* [13].

Results

Between 18 May 2011 and 23 January 2012, 20 MRSA colonized or infected neonates (including three sets of twins) were identified in the NICU (Fig. 2). Fifteen were outbreak cases, of whom four had MRSA infection: one each with purulent ear discharge and severe pneumonia and the two index cases, with fatal sepsis.

Case 1 had rapidly progressive sepsis and necrotizing pneumonia, with pneumatoceles. MRSA was isolated from tracheal aspirate and urine but not blood. Case 2 developed overwhelming sepsis, with MRSA in blood and CSF, 2 days after MRSA had been isolated from a nasal lesion. Both infants were treated empirically with vancomycin at the onset of symptoms and later with linezolid and/or clindamycin. Clinical presentations and outcomes of all 20 cases (see below) are summarized in Table 1.

In May 2011, surveillance cultures were collected from 138 NICU HCW; two (staff A and B) were colonized and successfully decolonized. In June 2011, a bundle of enhanced infection control measures was implemented: MRSA-




FIG. 2. Outbreak timeline between 18 May 2011 until 23 January 2012. PPE, personal protective equipment; HH, hand hygiene; HCW, healthcare worker.

TABLE 1. Clinical data for infants colonized or infected with methicillin-resistant *Staphylococcus aureus* (MRSA)

CASE	Sex/ethnicity	Gestational age, weeks	Birth weight, g	Birth	Underlying conditions (s)	Age in days at first isolation of MRSA	Risk factors	Clinical presentation	Infection/colonisation	Other manifestations	Site	Antimicrobial treatment	Outcome
1	F/Caucasian	26	730	LSCS	HMD, PDA, IUGR, XXX anomaly	9	UVC, CVC, ICC, TPN	Septic shock	Infection	Pneumothorax	ETT aspirate	Penicillin, gentamicin, vancomycin, cefotaxime, meropenem	Died
2	F/East or SE Asian	26	830	PROM, Vaginal breech	HMD, PDA, Grade 1 IVH	11	UVC, UAC, CVC, ICC	Septic shock, chest infection	Infection	Pneumothorax, anaemia of prematurity	Blood	Penicillin, gentamicin, meropenem, vancomycin	Died
3	F/Middle Eastern	28	1010	PROM, LSCS	HMD, ROP	2	CVC	Suspected sepsis	Colonisation	GORD	Surveillance swab	Penicillin, gentamicin, vancomycin	Survived
4	M/Middle Eastern	28	1310	PROM, LSCS	HMD, PDA, Grade 2 IVH	8	CVC	Suspected sepsis	Colonisation	GORD	Surveillance swab	Penicillin, gentamicin, vancomycin	Survived
5	M/Caucasian	40	4122	PROM, LSCS	None	8	PVC	R breast abscess	Infection	None	Umbilicus	Flucloxacillin, gentamicin, vancomycin	Survived
6	M/Caucasian	27	1210	LSCS	HMD, PDA	4	PVC	L purulent discharge from ear	Infection	CLD, GORD	Ear	Penicillin, gentamicin, vancomycin	Survived
7	M/Caucasian	25	780	LSCS	HMD, PDA, ROP	11	UVC, UAC, CVC	Sepsis, seizures, pneumonia	Infection	GORD, conjugated hyperbilirubinaemia	Eye	Penicillin, gentamicin, vancomycin, rifampicin, clindamycin, meropenem	Survived
8	M/South Asian	38	3100	PROM, NVB	None	2	PVC	Pustules on head	infection	None	Surveillance swab	Penicillin, gentamicin, flucloxacillin	Survived
9	M/Caucasian	38	4040	elective LSCS	None	3	None	Haematuria	colonisation	None	Bag urine	None	Survived
10	F/Caucasian	26	1000	LSCS	HMD, NEC, Grade 1 IVH	5	UVC, CVC	NEC	Colonisation	None	Surveillance swab	Penicillin, gentamicin, vancomycin, clindamycin	Survived
11	M/Caucasian	26	890	LSCS	HMD, PDA, ROP	141	UVC, UAC, CVC	Suspected sepsis	Colonisation	CLD, metabolic bone disease, bilateral inguinal hernias	Surveillance swab	Penicillin, gentamicin, cefotaxime, vancomycin, augmentin	Survived
12	F/Middle Eastern	27	1100	LSCS	HMD	1	CVC	Coagulopathy	Colonisation	GORD, PDA	Surveillance swab	Penicillin, gentamicin	Survived
13	F/Middle Eastern	27	1210	LSCS	HMD, PDA	1	UVC, UAC, CVC	Suspected sepsis	Colonisation	PDA	Surveillance swab	Penicillin, gentamicin	Survived
14	F/Caucasian	40	3195	NVB	HMD, PDA	10	None	L breast	Colonisation	None	Surveillance swab	Vancomycin gentamicin, vancomycin gentamicin	Survived
15	M	32	1260	LSCS	HMD, PDA	15	UAC, UVC, CVC	Suspected sepsis	Colonisation	CLD, Bilateral inguinal hernia, penoscrotal hypospadias	Surveillance swab	vancomycin gentamicin	Survived
16	M	32	1165	LSCS	HMD, PDA	15	UAC, UVC, CVC	Midgut volvulus	Colonisation	CLD, Bilateral inguinal hernia, penoscrotal hypospadias	Surveillance swab	Vancomycin gentamicin, Metronidazole	Survived
17	F	38	2640	Vaginal breech	Trisomy 21, AVSD	8	None	Heart failure	Colonisation	None	Surveillance swab	None	Survived
18	M/Middle Eastern	36 + 3	1820	NVB	HMD	1	PVC	Suspected sepsis	Colonisation	None	Surveillance swab	Penicillin, gentamicin	Survived
19	M/Middle Eastern	31 + 2	1890	NVB	HMD	9	PVC	Suspected sepsis	Colonisation	None	Surveillance swab	Penicillin, gentamicin	Survived
20	M/Caucasian	36 + 4	2420	LSCS	TTN	3	None	Asymptomatic	Colonisation	None	Surveillance swab	None	Survived

ASD, atrial septal defect; AVSD, atrioventricular septal defect; CLD, chronic lung disease; HMD, hyaline membrane disease; PDA, patent ductus arteriosus; GORD, gastroesophageal reflux disease; IVH, intraventricular haemorrhage; IUGR, intrauterine growth retardation; NEC, necrotising enterocolitis; NVB, normal vaginal birth; PROM, premature rupture of membranes; PVC, peripheral venous catheter; ROP, retinopathy of prematurity; UAC, umbilical arterial catheter, UVC, umbilical venous catheter; CVC, central venous catheter; ICC, intercostal catheter; TPN, total parenteral nutrition, TTN, Transient Tachypnoea of Newborn. Cases that isolated the outbreak strain are highlighted.

colonized infants were placed on contact precautions and cohorted until discharge; and a unit-wide review of hand hygiene practices, staff education on '5 Moments of Hand Hygiene' (Fig. 2) and enhanced environmental cleaning of the MRSA-colonized patient rooms were implemented.

Despite these measures, there was ongoing MRSA acquisition, including by three infants (twins and a singleton) who were positive on admission to the NICU, following Caesarean section delivery. Therefore, a second bundle of measures was implemented in September 2011. Additional or repeat MRSA screening of NICU staff was extended to staff in the delivery suite. Two of 146 NICU HCW (20 doctors, 107 nurses and 19 patient support assistants, technicians and cleaning staff) screened were colonized (staff C and D; both were negative in June). Two of 83 delivery suite staff (26 obstetricians, 33 midwives, 23 anesthetic staff and one other) screened were colonized (staff E and F). All colonized staff were successfully decolonized.

Accredited auditors performed daily '5 moments' hand hygiene audits and compliance improved from 81% in September to 95% in January 2012; audits were subsequently performed weekly. Patient zones were physically demarcated to improve adherence to contact precautions. Additional environmental screening yielded MRSA from three of 52 swabs: a blood gas machine, a milk trolley and a curtain. Five of 13 mothers (38%; 95% CI 31–45%) and both of two fathers screened were MRSA colonized (see Table 2).

Molecular typing results

RLB binary typing identified 18 indistinguishable MRSA isolates (the 'outbreak' strain) from 15 neonates, one HCW (staff C), one father and one arterial blood gas machine swab (Table 2). This strain was PVL positive, ST 22, *SCCmec* sub-type IV.3.1.2 (or IVc) and *spa* type t005. The PFGE patterns of these isolates were indistinguishable.

All colonized mothers carried non-outbreak strains, three of which matched those of their infants. The other two were mothers of outbreak cases (6 and 7). Staff C, who was colonized with the outbreak strain, had cared for outbreak cases 3 and 4. Staff D was colonized with the same non-outbreak strain as case 8, whom she had nursed. This multi-resistant strain was identified as ST772-V, *spa* type t657 (Bengal Bay clone). Six non-outbreak isolates with indistinguishable RLB types were isolated from a midwife (Staff F), three babies (cases 5, 9 and 14), one mother (9) and a delivery suite curtain. Apart from the mother/baby pair (case 9), there were no apparent epidemiological links; staff F was not involved in delivery of those babies. These strains were typical of STI-IV (USA400 or WA-MRSA-I), which is the second most common community-associated MRSA clone in Australia [14].

Discussion

We documented sustained nosocomial transmission of a virulent MRSA PVL-positive ST22-MRSA-IV in our NICU, affecting 15 neonates, one father and one HCW over a 7-month period. This is the first report of a nosocomial outbreak due to this clone in Australia. PVL, which is thought to be important in pathogenesis [15,16], probably contributed to fatal necrotizing pneumonia with pneumatocele formation, in one index case, and to septic shock and bullous lung abscesses in the other.

We demonstrated the utility of our novel mPCR/RLB typing method, to rapidly identify nosocomial transmission and define the outbreak by excluding MRSA-colonized infants carrying non-outbreak strains. Traditional MRSA typing methods are too expensive, too slow (PFGE, MLST) and/or have insufficient discriminatory power (MLST) for routine use. *spa* typing has been used, prospectively, to identify nosocomial transmission events [17] but is unsuitable in our institution, where the dominant MRSA clone (ST239-III) belongs to a single *spa* type (t037). Our binary typing method has a turnaround time of 12 h and high discriminatory power; it is inexpensive (~US\$2 per isolate) and up to 80 isolates can be typed in a single run, using two membranes. It allows targeted infection control interventions to be implemented promptly.

The index cases shared a room; it was not clear who was colonized first, but transmission between them, by HCW or fomites, was assumed. Their mothers were not screened; one may have acquired infection through vertical transmission or from an unidentified HCW. No common source for subsequent cases was found; it was unlikely to have been the HCW colonized with the outbreak strain, after a previously negative screen. The outbreak strain was isolated from only one of more than 50 environmental swabs, but demonstrates the potential for MRSA to be transmitted via fomites and the need for targeted environmental cleaning and enhanced hand hygiene. Twelve of 15 outbreak cases were delivered by Caesarean section, including three colonized within 24 h of birth, suggesting acquisition by some from a delivery suite source, as well as transmission within the NICU.

Further evidence favouring nosocomial, rather than vertical, transmission was that mothers of two outbreak cases were colonized with non-outbreak strains, whereas colonized mothers of babies with non-outbreak strains carried the corresponding strain. Several studies have suggested that maternal vaginal MRSA colonization leading to neonatal infection is uncommon [18–20].

TABLE 2. MRSA isolate information

Sample	Date	Antibiotic Resistances	PVL	MLST	SCCmec	spa	Binary typing result	(Predicted)*																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																									
								sea	sec	sed	pvl	N046	SAV0881	PV830RF2	SAV0858	phlI-4568	slcORD257	SA1801	SAV1974	slcORF182	ccrAB1	ccrC	mechI	E007	C0002	cadB																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																							
Baby 1	18/05/2011	Gen TmpSmz	+	22	IV.3.1.2	t005	37 248																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																										</

Antibiotic resistances are for non-beta lactam antibiotics only; antibiotics tested were chloramphenicol, ciprofloxacin (Cip), clindamycin (Clin), erythromycin (Ery), fosfomycin (Fos), fusidic acid, gentamicin (Gen), linezolid, mupirocin, nitrofurantoin, quinupristin-dalfopristin, rifampin and trimethoprim-sulphamethoxazole (TmpSmz). Each target in the binary typing system was interrogated using two probes (sense and antisense), black squares indicated positive probes. Isolates belonging to the outbreak strain are highlighted. MLST, SCCmec and spa results in bold represent actual results; those in italics are predicted based on the binary typing result.

Binary typing showed, and formal MLST, *spa* and *SCCmec* typing confirmed, that the outbreak strain was PVL positive and unrelated to the usual nosocomial clones seen in our institution (AUS2/3 [ST239 MRSA-III] and UK-EMRSA15 [ST22-MRSA IV]). PFGE confirmed that the outbreak isolates were indistinguishable. PVL-positive ST22 MRSA has been rarely identified, previously, in Australia [14].

ST22-MRSA-IV is generally synonymous with UK-EMRSA-15, which emerged in hospitals in the United Kingdom in 1991 [21], was subsequently identified in Germany where it has become the predominant MRSA clone [22] and has spread to hospitals and long-term care facilities, worldwide. It is characteristically ciprofloxacin resistant and gentamicin susceptible, may produce enterotoxin C but rarely PVL, harbours *SCCmec* subtype IVh and has *spa* type t022 or t032 (or a related variant) [23].

Our outbreak strain differs from EMRSA-15; it has a different *SCCmec* subtype (IV3.1.2 or IVc), *spa* type (t005), antibiogram (resistant to gentamicin and trimethoprim-sulphamethoxazole but susceptible to ciprofloxacin) and produces PVL, but not enterotoxin C. It has been reported in the United Kingdom [24,25] and Japan [26], where the index case reported recent travel to India. An outbreak of breast abscesses in postpartum women in Mumbai, India, was due to a similar MRSA clone, differing only by a single nucleotide polymorphism in one *spa* repeat [27]. There was no known link with the Indian subcontinent in our cases, but staff and patients at our institution come from a broad range of ethnic backgrounds, so recent importation of this clone remains a possibility.

Considering the large number of differences in the non-core genome, this clone probably arose from ST22-MSSA, independently of EMRSA-15 [24]. An outbreak of PVL-positive ST-22 MSSA with the same *spa* type and antibiogram has been reported from Italy, also amongst postpartum women and neonates [28].

The methods we used for MRSA screening included both PCR and broth enrichment to optimize turn-around time and sensitivity, followed by subculture. Previous studies suggest that PCR-based screening has high sensitivity but low reproducibility in neonates and that a combined approach, with confirmatory culture, is preferable [29]. For optimal neonatal screening, the combination of umbilical and nasal swabs has a reported sensitivity of >90% [30]. The screening protocol we used (pooled ear, nose, umbilicus and perineum) should have similar sensitivity. However, only nasal swabs were collected from staff, which may explain the major limitation of this investigation, namely our failure to identify a source, despite extensive screening of those involved in delivery and care of these infants.

Among the mothers of colonized infants screened for MRSA, the carriage rate (of non-outbreak strains) was high (38%). Previous reports suggest overall anovaginal MRSA carriage rates of 2–4%, with low rates of vertical transmission of MRSA infection to neonates [18,19]. Systematic screening is required to determine the true MRSA carriage rates in our antenatal population but was beyond the scope of this study.

Non-multiresistant MRSA is an emerging pathogen in the NICU, and can cause outbreaks with significant mortality. The virulent PVL-positive ST22-MRSA-IV strain that caused this outbreak was difficult to control, despite timely use of a novel typing system, which allowed accurate definition of the outbreak, and implementation of targeted infection control interventions that have, apparently, finally terminated the outbreak. At the time of writing there have been no new acquisitions of the outbreak strain for more than 4 months. EMRSA-15 is not the only ST22-MRSA-IV clone with epidemic potential.

Acknowledgements

We thank Francis Jenkins for assistance with RLB typing, Justin Ellem for laboratory assistance with MRSA surveillance, and staff of the Infection Prevention and Control Unit and the Centre for Newborn Care, Westmead Hospital. Molecular strain typing of MRSA was funded, in part, by a National Health and Medical Research (NHMRC) project grant #1010452 (CIA Gilbert).

Transparency Declaration

The authors declare no conflict of interests.

References

1. Healy CM, Hulten KG, Palazzi DL, Campbell JR, Baker CJ. Emergence of New Strains of Methicillin-Resistant *Staphylococcus aureus* in a Neonatal Intensive Care Unit. *Clin Infect Dis* 2004; 39: 1460–1466.
2. Heinrich N, Mueller A, Bartmann P, Simon A, Bierbaum G, Engelhart S. Successful management of an MRSA outbreak in a neonatal intensive care unit. *Eur J Clin Microbiol Infect Dis* 2011; 30: 909–913.
3. Saiman LMD, Cronquist ARN, Wu FMD et al. An Outbreak Of Methicillin-Resistant *Staphylococcus aureus* In A Neonatal Intensive Care Unit. *Infect Control Hosp Epidemiol* 2003; 24: 317–321.
4. Milstone AM, Goldner BW, Ross T, Shepard JW, Carroll KC, Perl TM. Methicillin-Resistant *Staphylococcus aureus* Colonization and Risk of Subsequent Infection in Critically Ill Children: importance of Preventing Nosocomial Methicillin-Resistant *Staphylococcus aureus* Transmission. *Clin Infect Dis* 2011; 53: 853–859.

5. Bannerman TL, Peacock SJ. *Staphylococcus, Micrococcus and Other Catalase-Positive Cocci*. In: Murray PR, Baron EJ, Jorgensen JH, Landry ML, Pfaller MA, eds, *Manual of Clinical Microbiology*, 9th edn. Washington, DC: American Society for Microbiology, 2007; 390–411.
6. Cai Y, Kong F, Wang Q et al. Comparison of single and multilocus sequence typing and toxin gene profiling for characterisation of methicillin resistant *Staphylococcus aureus* (MRSA). *J Clin Microbiol* 2007; 45: 3302–3308.
7. O'Sullivan MV, Kong F, Sintchenko V, Gilbert GL. Rapid identification of methicillin-resistant *Staphylococcus aureus* transmission in hospitals by use of phage-derived open reading frame typing enhanced by multiplex PCR and reverse line blot assay. *J Clin Microbiol* 2010; 48: 2741–2748.
8. Cai L, Kong F, Wang Q et al. A new multiplex PCR-based reverse line-blot hybridization (mPCR/RLB) assay for rapid staphylococcal cassette chromosome mec (SCCmec) typing. *J Med Microbiol* 2009; 58: 1045–1057.
9. O'Sullivan MV, Zhou F, Sintchenko V, Kong F, Gilbert GL. Multiplex PCR and reverse line blot hybridization assay (mPCR/RLB). *J Vis Exp* 2011; 49: pii: 2781.
10. Severiano A, Pinto FR, Ramirez M, Carriço JA. Adjusted Wallace Coefficient as a Measure of Congruence between Typing Methods. *J Clin Microbiol* 2011; 49: 3997–4000.
11. Murchan S, Kaufmann ME, Deplano A et al. Harmonization of pulsed-field gel electrophoresis protocols for epidemiological typing of strains of methicillin-resistant *Staphylococcus aureus*: a single approach developed by consensus in 10 European laboratories and its application for tracing the spread of related strains. *J Clin Microbiol* 2003; 41: 1574–1585.
12. Enright MC, Day NP, Davies CE, Peacock SJ, Spratt BG. Multilocus sequence typing for characterization of methicillin-resistant and methicillin-susceptible clones of *Staphylococcus aureus*. *J Clin Microbiol* 2000; 38: 1008–1015.
13. Chongtrakool P, Ito T, Ma XX et al. Staphylococcal cassette chromosome mec (SCCmec) typing of methicillin-resistant *Staphylococcus aureus* strains isolated in 11 Asian countries: a proposal for a new nomenclature for SCCmec elements. *Antimicrob Agents Chemother* 2006; 50: 1001–1012.
14. Coombs GW, Pearson J, Christiansen K, Nimmo GR. *Staphylococcus aureus* Programme 2010 (SAP 2010) Community Survey: MRSA Epidemiology and Typing Report. Available at: <http://www.agargroup.org/files/FED%20REPORT%20SAP210%20MRSA%20FINAL%20shrink.pdf> (last accessed 15 June 2012).
15. McAdams RM, Mazuchowski E, Ellis MW, Rajnik M. Necrotizing Staphylococcal Pneumonia in a Neonate. *J Perinatol* 2005; 25: 677–679.
16. Yee-Guardino S, Kumar D, Abughali N, Tuohy M, Hall GS, Kumar ML. Recognition and treatment of neonatal community-associated MRSA pneumonia and bacteremia. *Pediatr Pulmonol* 2008; 43: 203–205.
17. Mellmann A, Friedrich AW, Rosenkötter N et al. Automated DNA Sequence-Based Early Warning System for the Detection of Methicillin-Resistant *Staphylococcus aureus* Outbreaks. *PLoS Med* 2006; 3: e33.
18. Andrews WW, Schelonka R, Waites K, Stamm A, Cliver SP, Moser S. Genital tract methicillin-resistant *Staphylococcus aureus*: risk of vertical transmission in pregnant women. *Obstet Gynecol* 2008; 111: 113–118.
19. Beigi R, Hanrahan J. *Staphylococcus aureus* and MRSA colonization rates among gravidas admitted to labor and delivery: a pilot study. *Infect Dis Obstet Gynecol* 2007; 2007: 70876.
20. Pinter DM, Mandel J, Hulten KG, Minkoff H, Tosi MF. Maternal-infant perinatal transmission of methicillin-resistant and methicillin-sensitive *Staphylococcus aureus*. *Am J Perinatol* 2009; 26: 145–151.
21. Methicillin-resistant *Staphylococcus aureus* (MRSA). *Commun Dis Rep CDR Wkly* 1992; 2:33.
22. Witte W, Enright M, Schmitz FJ, Cuny C, Bräulke C, Heuck D. Characteristics of a new epidemic MRSA in Germany ancestral to United Kingdom EMRSA 15. *Int J Med Microbiol* 2001; 290:677–682.
23. Ghebremedhin B, König W, Witte W, Hardy KJ, Hawkey PM, König B. Subtyping of ST22-MRSA-IV (Barnim epidemic MRSA strain) at a university clinic in Germany from 2002 to 2005. *J Med Microbiol* 2007; 56:365–375.
24. Boakes E, Kearns AM, Ganner M et al. Molecular diversity within clonal complex 22 methicillin-resistant *Staphylococcus aureus* encoding Panton–Valentine leukocidin in England and Wales. *Clin Microbiol Infect* 2011; 17: 140–145.
25. Ellington MJ, Perry C, Ganner M et al. Clinical and molecular epidemiology of ciprofloxacin-susceptible MRSA encoding PVL in England and Wales. *Eur J Clin Microbiol Infect Dis* 2009; 28: 1113–1121.
26. Yamamoto T, Takano T, Yabe S et al. Super-sticky familial infections caused by Panton–Valentine leukocidin-positive ST22 community-acquired methicillin-resistant *Staphylococcus aureus* in Japan. *J Infect Chemother* 2012; 18: 187–198: 1–12.
27. Manoharan A, Zhang L, Poojary A, Bhandarkar L, Koppikar G, Robinson DA. An outbreak of post-partum breast abscesses in Mumbai, India caused by ST22-MRSA-IV: genetic characteristics and epidemiological implications. *Epidemiol Infect* 2012; 1–4. [Epub ahead of print].
28. Tinelli M, Monaco M, Vimercati M, Ceraminiello A, Pantosti A. Methicillin-susceptible *Staphylococcus aureus* in skin and soft tissue infections, Northern Italy. *Emerg Infect Dis* 2009; 15: 250–257.
29. Sarda VMD, Molloy AMSN, Kadkol SMD, Janda William MP, Hershov RMD, McGuinn MMD. Active Surveillance for Methicillin Resistant *Staphylococcus aureus* in the Neonatal Intensive Care Unit. *Infect Control Hosp Epidemiol* 2009; 30: 854–860.
30. Rosenthal A, White D, Churilla S, Brodie S, Katz KC. Optimal Surveillance Culture Sites for Detection of Methicillin-Resistant *Staphylococcus aureus* in Newborns. *J Clin Microbiol* 2006; 44: 4234–4236.